

Isolation and molecular identification of *Serratia nematodiphila* associated with Red Palm Weevil, *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae) as bio-insecticide in Egypt

Mahmoud Abbas Ali¹, Mervat A. B. Mahmoud², Muhammad Shoaib³, Zeeshan Ahmad Bhutta^{4*}, Nada M. Ali⁵, Nadeem Ali⁶, Hani Z. Asfour⁷, Nisreen Rajeh⁸, Mohamed R. Eletmany^{9,10}

¹Department of Plant Protection, Faculty of Agriculture, South Valley University, Qena, Egypt

²Department of Zoology, Faculty of Science, South Valley University, Qena, Egypt

³Key Laboratory of New Animal Drug Project, Gansu Province/Key Laboratory of Veterinary Pharmaceutical Development, Ministry of Agriculture and Rural Affairs/Lanzhou Institute of Husbandry and Pharmaceutical Sciences of the Chinese Academy of Agricultural Sciences, Lanzhou 730050, China

⁴Laboratory of Veterinary Immunology and Biochemistry, College of Veterinary Medicine, Chungbuk National University, Cheongju 28644, Republic of Korea

⁵Department of Chemistry, Faculty of Science, Al-Baha University, Al-Baha 65799, Saudi Arabia

⁶Center of Excellence in Environmental Studies, King Abdulaziz University, Jeddah, 21589, Saudi Arabia

⁷Department of Medical Microbiology and Parasitology, Faculty of Medicine, King Abdulaziz University, Jeddah 21589, Saudi Arabia

⁸Department of Clinical Anatomy, Faculty of Medicine, King Abdulaziz University, Jeddah, 21589, Saudi Arabia

⁹Department of Chemistry, Faculty of Science, South Valley University, Qena, Egypt

¹⁰TECS Department, Wilson College of Textiles, NC State University, Raleigh 27606, USA

Received:

December 09, 2023

Accepted:

December 28, 2023

Published Online:

January 19, 2024

Abstract

The red palm weevil (RPW), *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae) is a major date palm pest. In this study, we aimed to isolate and identify the *Serratia nematodiphila* from RPW as potential biocontrol agents. We isolated the bacteria from infected RPW larvae and adults and identified using colony morphology characteristics, biochemical tests, and PCR followed by 16S rRNA sequencing. This is the first study reporting the *Serratia nematodiphila* as an extracellular symbiont of RPW from Egypt. The potential of this bacteria to be used as biocontrol agent was conducted by a screening bioassay through its effect on RPW eggs. The study noted that treated eggs were unable to hatch and not turned red in color, indicating the potential of this bacteria to be used as bio-pesticide. These results presented novel insights into the microbiome of RPW and suggest the potential of *Serratia nematodiphila* as a biocontrol agent for RPW management. Moreover, further studies are required to explore the mechanism and potential of these bacteria in field applications. Nevertheless, this study provides a promising direction for the development of sustainable and environmentally friendly RPW management strategies.

Keywords: Red Palm Weevils (RPW), *Rhynchophorus ferrugineus*, Coleoptera, Curculionidae, *Serratia nematodiphila*, Biocontrol agent

How to cite this:

Ali MA, Mahmoud MAB, Shoaib M, Bhutta ZA, Ali NM, Ali N, Asfour HZ, Rajeh N and Eletmany MR. Isolation and molecular identification of *Serratia nematodiphila* associated with Red Palm Weevil, *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae) as bio-insecticide in Egypt. Asian J. Agric. Biol. 2024(3): 2023352. DOI: <https://doi.org/10.35495/ajab.2023.352>

*Corresponding author email:
zee@chungbuk.ac.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License. (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Introduction

The date palm, *Phoenix dactylifera* L. belong to *Arecaeae* family is a fruit crop of tropical regions. Its cultivation has spread all over the world which is also known as oldest fruit tree planted in Middle Eastern countries, especially in the Arab Gulf states and Egypt. The Arab countries dominate the world's date production, while Egypt is the country that majorly producing dates worldwide. Egypt produces more than one million metric tons of fresh dates annually which accounts almost 18 % of all dates produced globally and 23 % of Arab production, with 1.590.414 million tons of the total estimated global production of nine million tons in 2017 (El-Sharabasy and Rizk, 2019). Date palms are susceptible to many destructive pests, which negatively affect production as well as quality (Alatawi et al., 2018).

The RPW, *Rhynchophorus ferrugineus* (Olivier), is an important date palm pest around the world (Al-Ayedh, 2020; Munawar et al., 2020). Since mid-1980s, RPW known to be disseminated at large geographical area including Asia, America, Africa and Europe (EPPO, 2020). From its origin, it spread widely through the trade of palm trees and offshoots from infected areas. The larvae is the main destruction stage. It drills into palm trees and eats succulents, remain hidden for most of its life cycle. In 2009, the expected losses in Gulf Cooperation Council countries of the Middle East were estimated to be 5.18 to 25.92 million USD due to the severe infestation of date palms at a rate of 1.0% to 5.0% respectively (El-Sabea et al., 2009). The Food and Agriculture Organization of United Nations classified *R. ferrugineus* as a class-1 pest in the Middle East and North Africa (MENA) regions (El-Shafie and Faleiro, 2020) and A2 pest in the Europe (Faleiro et al., 2019). It poses a significant problem to the livelihood of date palm producers in rural areas of MENA.

There are various strategies that can be applied to control red palm weevil (RPW), including chemical, cultural, biological, and physical control methods. The chemical control comprises the use of insecticides, such as chlorpyrifos and deltamethrin, to kill RPW larvae and adults. However, cultural control involves various implementing measures such as pruning, sanitation, and trapping to reduce RPW infestations while physical prevention consists of practicing the physical barriers, for example sticky bands, to restrict RPW from climbing up the palm trees. The biological way of control uses the natural enemies, such as

predators and parasites, to control RPW population. The use of all these control methods called as integrated pest management (IPM) which is a comprehensive approach to manage RPW populations sustainably. The selection of an appropriate control strategy depends on various factors, such as severity of RPW infestation, local environment, and economics.

Biological control is known as one of the best strategies in combating the red palm weevil because its less expensive, ecofriendly, minimum legal and public health concerns (Al-Dosary et al., 2016; Ramzan et al., 2023). As the biological control uses the vital live enemies which can be pathogens, predators, and parasites such as bacteria and fungi (Zhu et al., 2010; Pu and Hou, 2016; Abd-Elfatah et al., 2023). It provides advantageous conditions for the environment and economy since yield loss may be reduced without causing unfavorable natural effects, which can happen when pesticides are used (Bianchi et al., 2006). Entomopathogenic bacteria belong to the major microbial families which are *Bacillaceae*, *Streptococcaceae*, *Micrococcaceae*, *Enterobacteriaceae*, and *Pseudomonadaceae* (Kaya et al., 1993; Mehmood and Ashraf, 2023). Previously, *B. thuringiensis* subspecies *kurstaki* was isolated from RPW larvae in Egypt (Alfazariy, 2004), *Serratia marcescens* and *Serratia nematodiphila* in China (Zhang et al., 2009; Pu and Hou, 2016) was used to successfully control the RPW under laboratory settings. This is the first study isolated and identified the *Serratia nematodiphila* symbiotically associated with RPW, *Rhynchophorus ferrugineus* (Olivier) from Egypt and were assessed for bio-insecticide activity against *Rhynchophorus ferrugineus*.

Material and Methods

Red palm weevil collection and laboratory culture

RPW larvae, pupae, and adults were collected from infested date palms (*Phoenix dactylifera* L.) located in various areas of Qena governorate, namely, South Valley Agricultural farm, Qeft, and Qus, from April 2020 to August 2021. Adult RPW were reared in plastic boxes (20×15×10 cm) fitted with a perforated cover for ventilation. It was provided with pieces of peeled Sugarcane reeds for feeding and laying eggs to obtain the eggs and the following stages, such as larvae and pupae. The rearing was carried out under laboratory conditions at 25 °C and 65% HR. The larvae were provided with reed sticks that were



hollowed out for feeding, and it was possible to get the pupae, which were placed in the same boxes as before until the adult weevil emergence.

Isolation of bacteria

The bacteria were isolated from already collected RPW pupae from different geographical location. Two knots were taken from the bacteria inside the cocoon, and they were dissolved in about 5 ml of sterile distilled water. The amount of a knot was taken from it using a sterile inoculation loop, and streaked on nutrient agar plates (10 g peptone: 3 g meat extract, 5g NaCl, 15 g agar and 1000 ml distilled, pH 7.3). The plates were incubated in an incubator at a temperature of 29°C for 48 hours. Different single isolated colonies were selected and re-cultured for further purification and to observe colony characteristics following the same incubation conditions. They were re-cultured again to ensure the purity of the bacterial cultures. The resultant growth of different colonies was observed for individual characteristics such as colony texture, color, size, and transparency. The identified strains were stored first at 4°C and then at -80 °C in 25% glycerol for further processing.

16S ribosomal DNA (rDNA) based molecular identification

Extraction of bacterial DNA

The bacterial DNA was extracted in Microbial Genetics Department, Biotechnology Research Institute, National Research Centre, Cairo, Egypt from the fresh overnight grown bacterial culture using Qiagen genomic DNA kit (Qiagen, Shanghai, China) as per guidelines of manufacturer. The extracted DNA was stored at -20 °C.

Identification by PCR amplification of the 16S rDNA gene

All extracted DNA was subjected to PCR amplification of 16S rDNA gene and later on sequencing. The PCR amplification of 16S rDNA gene was done using the universal primer as described earlier 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-CTACGGCTACCTTGTTCAG-3'). The PCR solution (50µL) consists of 4 µL DNA as template, 2 µL of each 10mM forward and reverse primers, 25 µL of Master Mix and 17 µL of nuclease free water. The PCR were run under following conditions; first denaturation at 94 °C for 3 minutes along with 35 cycles of denaturation at 94 °C for 30

sec, annealing at 55 °C for 30 sec, and initial extension at 72 °C for 1 min with a final extension at 74°C for 10 min and stored at 4 °C for infinite time. The PCR product yielded were analyzed on 1.0% (wt./vol) agarose gel after staining with ethidium bromide.

Agarose gel electrophoresis, purification, and sequencing

Agarose powder was dissolved in 1X TBE buffer at 1% and boiled in hot air oven for few minutes, then the ethidium bromide, gel staining dye was mixed in pre-heated gel solution when the temperature reaches approximately to 55°C. Then the gel was poured in the plastic tray and the comb was placed immediately and removed after solidification. Then the gel was placed in 1X TBE electrophoresis buffer filled in electrophoresis apparatus. The whole PCR product were loaded in each well and compared with 5 µl of 2500 bp DNA ladder. Later, the PCR product was extracted and purified from agarose gel using Wizard Genomic DNA Purification Kit (Promega, USA). The purified PCR product were sent for sequencing by HVD Life Science, Germany.

Phylogenetic analysis

Chromatograms were edited and trimmed using Bio Edit software. Edited sequences were aligned using Clustal X of Clustal W packages. The 16S rDNA sequence obtained were compared with highly similar publicly available sequences in NCBI database. All of the sequences were aligned and phylogenetic tree was constructed by using neighbor-joining (NJ) method in MEGA 11 (Kumar et al., 2018).

Bio-assay to check the insecticide activity on RPW eggs

The bacterial culture was mixed in sterile water and diluted into three different concentrations; 1×10^6 , 1.5×10^6 , 1×10^7 CFU/mL, which was further used in the whole bioassay. Sterile water was used as control group and the three bacterial concentrations were used as three treatments on RPW eggs. A total of 30 eggs were evaluated for each treatment and repeated thrice. Treatments entailed uniformly dosing 10 mL of the solution under test over the dish and its contents before adding 10 nondestructive, freshly laid eggs to a Petri dish (90 mm) with cotton on the bottom. Normally, it takes 3-5 days for RPW eggs to hatch, we counted the number of hatching larvae every day up to the seventh day. The larvae that had just hatched were swiftly



transferred to other petri dish to protect the other eggs from harm. The eggs were deemed non-viable if they did not hatch after seven days. By comparing the quantity of viable eggs before and after the administration of treatments, egg hatching rates were calculated.

Statistical analysis

The statistical analyses were performed using SPSS computer program (SPSS Inc., Chicago, IL, USA) at $\alpha=0.05$ (level of significance). The data of the egg hatching rate, adjusted mortality, and boring rate at the various dosages were compared using one-way ANOVA. Tukey's honestly significant difference (HSD) test was applied to compared the means within and between the groups.

Results

Isolation and colony characteristics of *Serratia nematodiphila*

Firstly, RPW larvae were detected by properties such as viscous layer on their surface and foul smelling. The surface was disinfected and washed with sterile Normal saline (0.9%) and isolation of bacteria was done on nutrient agar. The bacteria showed creamy color colonies, with smooth texture, bulging shape,

irregular margins and transparent which are indicative of particular bacteria.

16S rDNA gene-based identification and sequence analysis

The 16S rDNA gene was amplified by PCR to identify the bacterial isolates. The PCR product was pictured on 1% agarose gel at almost 1400-1500 bp length. The amplified gene was sequenced using Sanger sequencing technique and it showed 100% similarity to already submitted sequence in NCBI Genbank under the accession number, NR_044385.1. The bacterial sequence was blasted in NCBI blast to query the related sequences in NCBI database to identify clonal relationship (Table 1). Chromatograms were edited and trimmed using Bio Edit software (Fig. 1). The phylogenetic tree was constructed using Neighbor-Joining method in MEGA-11 by aligning the sequences through Clustal W (Fig. 2). The phylogenetic tree showed that the isolated strain was belonging to different clade as compared to similar species under the same genus but there was a high degree of similarity in their genomes ($\geq 95\%$). Apart from this, the sequence was also greatly similar, 95.6% to *Pectobacterium spp.*, *Coronobacter dublinensis subspp.* *Lausanensis*, *Citrobacter murlinae*, and *Enterobacter Kobei* strains.

Table 1. BLAST result of 16S rDNA gene sequence of *Serratia nematodiphila*

Bacterial Strains	Max Score	Total Score	Query Cover	E value	Percent Identity	Accession
<i>Serratia nematodiphila</i>	1589	1589	90%	0.0	98.03%	NR_044385.1
<i>Serratia marcescens</i> subsp. <i>marcescens</i> , JCM 1239	1587	1587	90%	0.0	97.93%	NR_113236.1
<i>Serratia marcescens</i> subsp. <i>marcescens</i> , DSM 30121	1578	1578	90%	0.0	97.82%	NR_041980.1
<i>Serratia surfactantfaciens</i> , YD25	1567	1567	90%	0.0	97.60%	NR_169468.1
<i>Serratia ureilytica</i> , NiVa 51	1533	1533	90%	0.0	96.94%	NR_042356.1
<i>Serratia entomophila</i> , DSM 12358	1478	1478	90%	0.0	95.86%	NR_025338.1
<i>Serratia ficaria</i> strain DSM 4569 16S ribosomal RNA, partial sequence	1472	1472	90%	0.0	95.75%	NR_041979.1
<i>Serratia ficaria</i> , NBRC 102596	1469	1469	90%	0.0	95.64%	NR_114155.1
<i>Serratia ficaria</i> , JCM1241	1467	1467	90%	0.0	95.64%	NR_112005.1
<i>Pectobacterium aroidearum</i> , SCRI 109	1467	1467	90%	0.0	95.63%	NR_159926.1
<i>Serratia odorifera</i> , PADG 1073	1461	1461	90%	0.0	95.52%	NR_037110.1






Figure-1. Chromatogram of the 16S rDNA gene sequence of isolated *Serratia nematodiphila*



Figure-2. Phylogenetic tree based on partial 16S rDNA sequences retrieved from NCBI database, showing the association between *Serratia nematodiphila* and other species.

Bioassay to check the insecticide activity

The effect of *Serratia nematodiphila* on RPW eggs was observed, as soon as the bacterial concentrations applied, red color appeared on eggs after 3 days (Fig. 3b). After application of different bacterial isolate concentrations sprayed on RPW eggs, hatching rates were significantly affected ($F_{3,8} = 161.222$, $P = 0.000$, Table 2). As the bacterial concentration increased, there was a tendency for fewer eggs to hatch (Table 3). The egg's average hatching rate were ranged between 3.3 and 8.67 which was considered as significant difference ($p < 0.05$). The eggs with the highest response rate to effectively dying were with 1×10^7 CFU/mL concentration. In contrast, there was a noticeable difference between the other various effective concentrations (Fig. 4A). Similar pattern was observed against larvae (Fig. 4B).




Figure-3. RPW eggs (A) Control group 0X, (B) treated eggs after 3 days 40X, (C) treated eggs after 5 days 40X.

Table-2. One-Way ANOVA test results

Treatments	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	120.917	3	40.306	161.222	0.000
Within Groups	2.000	8	0.250	-	-
Total	122.917	11	-	-	-

Determination of LC_{50} and LC_{90} of different biological agents against RPW eggs and larvae

The results of present study revealed the susceptibility of different biological agents to red palm weevil eggs and larvae, emphasizing key parameters such as LC_{50} , Index, Resistance Ratio (RR), Slope with uncertainty, and LC_{90} . The LC_{50} of *Serratia nematodiphila* egg showed that it was more susceptible to the treatment compared to other microorganisms listed in Table 3. This suggests that *Serratia nematodiphila* may have fewer defense mechanisms or a higher inherent vulnerability to the treatment compared to the other species. On the other hand, *Aspergillus flavus*, noted to have a more robust defense mechanism, a slower metabolic rate, or a lower uptake of the treatment. The other species, *Metarhizium anisopliae*, *Beauveria bassiana*, and *Trichoderma harzianum* exhibited varying degrees of susceptibility between these two species. Overall, the study provides insight into the varying responses of different species to the treatment (Table 3).




Figure-4. Effect of different concentrations of different biological agents on the (A) eggs and (B) larvae of RPW; Different superscripts on bar graphs indicate significant difference ($p < 0.05$) within the group while asterisk (***) indicate significant difference between the groups.

Table- 3. LC₅₀ and LC₉₀ of different biological agents against RPW eggs

Biological agents	LC ₅₀	Index	RR	Slope	Slope +/-	LC	LC ₉₀
<i>Serratia nematodiphila</i>	253770	100	1	0.722	0.159	2.54E+05	1.51E+07
<i>Metarhizium anisopliae</i>	3266100	7.77	12.87	0.859	0.146	3.27E+06	1.01E+08
<i>Beauveria bassiana</i>	3931000	6.456	15.49	1.052	0.149	3.93E+06	6.50E+07
<i>Trichoderma harzianum</i>	4937200	5.14	19.455	0.953	0.149	4.94E+06	1.09E+08
<i>Aspergillus flavus</i>	5138300	4.939	20.248	1.207	0.155	5.14E+06	5.92E+07

LC₅₀= LC denotes lethal concentration of any drug or treatment which will results in 50% death of specie/insect/animal while LC₉₀ indicates 90% death rate. RR denotes the resistance ratio which indicate the resistance exhibited by microorganism to a specific drug or biological agent.

Table-4. LC₅₀ and LC₉₀ of different biological agents against RPW larvae

Biological agents	LC ₅₀	Index	RR	Slope +/-	LC ₉₀
<i>Metarhizium anisopliae</i>	330350	119.261	0.838	0.491±0.147	1.35E+08
<i>Serratia nematodiphila</i>	393980	100	1	0.796±0.157	1.60E+07
<i>Trichoderma harzianum</i>	774210	50.888	1.965	0.76±0.149	3.77E+07
<i>Aspergillus flavus</i>	1826800	21.567	4.637	0.569±0.142	3.26E+08
<i>Beauveria bassiana</i>	2973800	13.248	7.548	0.337±0.155	1.89E+10

LC₅₀= LC denotes lethal concentration of any drug or treatment which will results in 50% death of specie/insect/animal while LC₉₀ indicates 90% death rate. RR denotes the resistance ratio which indicate the resistance exhibited by microorganism to a specific drug or biological agent.

Notably, *Metarhizium anisopliae* exhibited a significantly lower LC₅₀ (330350) with a high Index (119.261), suggesting superior efficacy. *Trichoderma harzianum* showed a moderate Index (50.888) and RR of 1.965, indicating reasonable effectiveness. *Aspergillus flavus* and *Beauveria bassiana* demonstrated higher LC₅₀ values and substantial RR, implying reduced potency against red palm weevil larvae (Table 4). These findings underscore the importance of assessing multiple parameters for a comprehensive evaluation of biological agents in pest management strategies.

Discussion

Previously, a great attention has been applied to control agriculture pests using various microbes such as fungi (Zhong and Chen, 2012; Buentello-Wong et al., 2015). Few researches have been conducted to evaluate the role of bacterial agents as biocontrol against RPW. To our knowledge, this is the first study which confirm the presence of the external symbiotic bacteria, *Serratia nematodiphila* specie associated with RPW. The genus *Serratia* includes almost 10 species and they are well known as red-pigmented bacteria because of a chemical called as Prodigiosin. Some of its species have a potential entomopathogenic role in control of pests such as *Serratia marcescens*

(Singh et al., 2008). Moreover, traditional biocontrol microbes include and limited to *Metarhizium*, *Bacillus*, *Beauveria*, nematodes and polyhedrosis viruses. The bacterial strain, *Serratia nematodiphila* will provide a new research horizon to be used as biocontrol agent against RPW in Egypt to eradicate *Rhynchophorus ferrugineus* (RPW).

The bacterial strain was isolated and identified successfully through microbiological and molecular techniques such as PCR and sequencing. The phylogenetic tree showed a close relationship with other *Serratia* species as shown in Fig 2. Moreover, the bioassay demonstrated the effect of this bacterial strain at different concentrations on the hatching of RPW eggs. Our findings demonstrated that the uppermost concentration (1×10^7 CFU/mL) decreased the hatching of eggs by 13.3%. As compared to the findings of previous researches, our findings proved that the *Serratia nematodiphila* can control the hatching the eggs which will avoid the date palm infestation from RPW larvae. Most previous studies concentrated on isolation, identification, but few studies focused on *Serratia* entomopathogenic effect on insects such as *S. marcescens* cell-free culture supernatant effected anti-feeding activity significantly and mortality to *P. Blanchard* (Pineda-Castellanos et al., 2015). *Serratia marcescens* causes the insect death. Previously, it was confirmed that *S. marcescens*



had significant insecticidal effect on the larvae of *S. exigua* and causes high mortality (Konecka et al., 2019). Another study conducted by isolated *S. marcescens* and *Serratia nematodiphila* from mangoes also present its anti-fungal activity (Trejo-López et al., 2022) and its reported previously that *Serratia* species produce a chemical compound known as prodigiosin which has good antimicrobial activity (Darshan and Manonmani, 2015; Lapenda et al., 2015).

In present study, five different microbial species were tested for their entomopathogenic effect on RPW eggs hatching rate and larvae death. It was noted that *Serratia nematodiphila* exhibited higher antimicrobial activity in terms of eggs hatching and larvae deaths as compared to other microbial species. Among the other microbial species, *Beauveria bassiana* exhibited higher antimicrobial activity as compared to *Metarhizium anisopliae* > *Trichoderma harzianum* > *Aspergillus flavus* on the hatchability of eggs. These results are in consistent with the findings of El Hussein (2019) that *Beauveria bassiana* causes the high mortality of RPW at different stages of life cycle. Another study conducted by Cito et al. (2014) studied the biochemical and antimicrobial activities of two *Metarhizium* strains, *Metarhizium anisopliae* and *Metarhizium pingshaense*. Both of the strain showed entomopathogenic activity by producing toxins and degrading proteases. Similarly, the pathogenic activity of *Trichoderma harzianum* as biocontrol agent has also been studied against other pathogens (Alahmadi et al., 2012; Sundaravadivelan and Padmanabhan, 2014) but not too much studies are reported. While in case of larvae deaths, the antimicrobial efficacy was in following order; *Trichoderma harzianum* > *Metarhizium anisopliae* > *Beauveria bassiana* > *Aspergillus flavus*. Among all these species, *Aspergillus flavus* showed lowest antimicrobial activity. *Aspergillus flavus* can also be the potential candidate for biological control of RPW as studied previously in Hail, Northern Saudi Arabia (Alanazi et al., 2020). These findings suggest *Serratia nematodiphila* as potential candidate for biological control of RPW in Egypt.

However, the bioassay was conducted in strictly controlled laboratory conditions. The variation between current study findings may be due to different bacterial strains, different environmental conditions, and restricted research on current bacterial strains worldwide. Therefore, more studies are needed evaluate the bio-insecticidal efficacy of *S. nematodiphila* in field trials in addition to investigate

the toxicity and applications to develop a solid basis for future utilization and development at large scale. Another concern which need to be studied in future is to evaluate the toxicity effect on non-target species such as humans and animals. It is of much concern to check their effects on non-target species to develop highly specific, less toxic and efficient microbial agents not only to control RPW but also other invasive species such as coleopteran pests.

A limitation of the current study is that the bioassays demonstrating insecticidal efficacy of *Serratia nematodiphila* were conducted under controlled laboratory conditions. While the results are promising, further studies are needed to evaluate the biocontrol potential of this bacterium under real-world field conditions against RPW infestations in date palm plantations. Additional factors like environmental influences, application methods, and persistence in the field may impact the insecticidal activity observed in this lab investigation. Field trials will be important to validate the potential of *Serratia nematodiphila* as a viable biopesticide for managing RPW in agriculture. This could be highlighted as an area for future investigation. Identification of bacterial diversity that are pathogenic to insects can lead to development of new biocontrol agents and our research reports highlighted the role of newly identified bacteria against RPW with high pesticide activity which make it a as future candidate for further research as a new bio-insecticide and recommends field studies.

Conclusion

The protracted utilization of chemical pesticides is discouraged due to its potential harm to beneficial insects, contribution to environmental pollution, and the development of resistance in *R. ferrugineus* populations, rendering pest management more challenging. Consequently, there is an urgent need for alternative and integrated pest management approaches that effectively addresses the *R. ferrugineus* infestation along with environmental safety. Such approaches may encompass biological control strategies, such as the deployment of natural predators or the utilization of pheromone traps to disrupt the mating patterns. Moreover, raising awareness among the garden managers and general population regarding the significance of early detection and preventative measures is pivotal in curtailing the further spread of this invasive species.



Further investigation is necessary to better comprehend the extent of its impact.

Acknowledgment

The authors would like to thank South Valley University for the assistance.

Disclaimer: None.

Conflict of Interest: None.

Source of Funding: The authors would like to thank South Valley University and the Egyptian Cultural Educational Bureau in Washington DC (GM1101)

Contribution of Authors

Ali MA: Conceptualization, methodology and original draft preparation

Mahmoud MAB: Project administration, data curation and software input

Shoaib M, Bhutta ZA & Rajeh N: Visualization, writing, reviewing and editing of manuscript

Ali NM & Asfour HZ: Formal analysis, writing, reviewing and editing of manuscript

Ali N & Eletmany MR: Funding acquisition, writing, reviewing and editing of manuscript

References

- Abd-Elfatah EB, Fawzi EM, Elsheikh HA and Shehata AA, 2023. Prevalence, virulence genes and antibiogram susceptibility pattern of *Staphylococcus aureus* and *Streptococcus agalactiae* isolated from mastitic ewes. *Int. J. Vet. Sci.* 12(2):152-160.
- Al-Ayedh H, 2020. Red palm weevil biology. *Red Palm Weevil: Guidelines on Management Practices*. Rome: FAO
- Al-Dosary NM, Al-Dobai S and Faleiro JR, 2016. Review on the management of red palm weevil *Rhynchophorus ferrugineus* Olivier in date palm *Phoenix dactylifera* L. *Emir. J. Food Agric.* 28(1):34-44.
- Alahmadi SS, Ouf SA, Ibrahim RA and El-Shaikh KA, 2012. Possible Control of Date Palm Stag Beetle, *Lucanus cervus* (L.)(Coleoptera: Lucanidae), Using Gut Protease Inhibitors of Different Bio-control Agents. *Egypt. J. Biol. Pest Control.* 22(2):93-101.
- Alanazi NA, Sulieman A, Alshammari NI, Al-Azmi M, Aziz A, Elbadri GA and Stephenson SL, 2020. Fungi related with the red palm weevil (*Rhynchophorus ferrugineus*) in the Hail Area, Northern Saudi Arabia. *Biosci. Biotech. Res. Commun.* 13(4):2233-2239.
- Alfazariy A, 2004. Notes on the survival capacity of two naturally occurring entomopathogens on the red palm weevil *Rhynchophorus ferrugineus* (Olivier) (Coleoptera: Curculionidae). *Egypt. J. Biol. Pest Control* 14:423.
- Bianchi FJ, Booij C and Tscharncke T, 2006. Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proc. Royal Soc. B Biol. Sci.* 273(1595):1715-1727.
- Buentello-Wong S, Galán-Wong L, Arévalo-Niño K, Almaguer-Cantú V and Rojas-Verde G, 2015. Characterization of Cry Proteins in Native Strains of *Bacillus thuringiensis* and Activity Against *Anastrepha ludens*1. *Southwest. Entomol.* 40(1):15-24.
- Cito A, Mazza G, Strangi A, Benvenuti C, Barzanti GP, Dreassi E, Turchetti T, Francardi V and Roversi PF, 2014. Characterization and comparison of *Metarhizium* strains isolated from *Rhynchophorus ferrugineus*. *FEMS Microbiol. Lett.* 355(2):108-115.
- Darshan N and Manonmani HK, 2015. Prodigiosin and its potential applications. *J. Food. Sci. Technol.* 52(9):5393-5407.
- El-Sabea AM, Faleiro J and Abo-El-Saad MM, 2009. The threat of red palm weevil *Rhynchophorus ferrugineus* to date plantations of the Gulf region in the Middle-East: an economic perspective. *Outlooks Pest Manage.* 20(3):131-134.
- El-Shafie HAF and Faleiro JR, 2020. Red palm weevil *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae): Global invasion, current management options, challenges and future prospects. *Invasive Species-Introduction Pathways, Economic Impact, and Possible Management Options*. pp. 1-30.
- El-Sharabasy S and Rizk R, 2019. Atlas of date palm in Egypt.
- El Hussein MM, 2019. Efficacy of the fungus *Beauveria bassiana* (Balsamo) Vuillemin on the red palm weevil *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae) larvae and adults under laboratory conditions. *Egypt. J. Biol. Pest Control.* 29(1):58.



- EPPO, 2020. *Rhynchophorus ferrugineus* (RHYCFE)[Datasheet] EPPO Global Database. URL: <https://gd.eppo.int/taxon/RHYCFE/datasheet>.
- Faleiro JR, Ferry M, Yaseen T and Al-Dobai S, 2019. Overview of the gaps, challenges and prospects of red palm weevil management. *Arab. J. Plant Prot.* 37(2):170-177
- Alatawi JF, Zain ul Abidin S, Mirza HJ and Kamran M, 2018. Functional response of the predatory mite *Cydnoseius negevi* (Swirski & Amitai) (Acari: Phytoseiidae) to the *Oligonychus afrasiaticus* (Mcgregor) and *Tetranychus urticae* Koch (Acari: Tetranychidae). *Asian J. Agric. Biol.* 6(2): 265-277.
- Kaya HK, Bedding RA and Akhurst RJ, 1993. An overview of insect-parasitic and entomopathogenic nematodes. *Nematodes and the biological control of insect pests*:1-10.
- Konecka E, Mokracka J, Krzysińska S and Kaznowski A, 2019. Evaluation of The Pathogenic Potential of Insecticidal Strains to Humans. *Pol. J. Microbiol.* 68(1):185-191.
- Kumar S, Stecher G, Li M, Knyaz C and Tamura K, 2018. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol.* 35(6):1547.
- Lapenda JC, Silva PA, Vicalvi MC, Sena KX and Nascimento SC, 2015. Antimicrobial activity of prodigiosin isolated from *Serratia marcescens* UFPEDA 398. *World J. Microbiol. Biotechnol.* 31(2):399-406.
- Mehmood S and Ashraf M, 2023. Antimicrobial resistance and virulence determinants of *E. coli* in bovine clinical mastitis in dairy farms. *Continental Vet. J.* 3(1):54-59.
- Munawar A, Ishaque Mastoi M, Sohail S and Bashir F, 2020. The compatibility of single and combined applications of the entomopathogenic nematode, *Heterorhabditis indica* with imidacloprid against red palm weevil, *Rhynchophorus ferrugineus* (Oliv.). *Asian J. Agric. Biol.* 8(3):315-322.
- Pineda-Castellanos ML, Rodríguez-Segura Z, Villalobos FJ, Hernández L, Lina L and Nuñez-Valdez ME, 2015. Pathogenicity of isolates of *Serratia marcescens* towards larvae of the scarab *Phyllophaga blanchardi* (Coleoptera). *Pathogens.* 4(2):210-228.
- Pu YC and Hou YM, 2016. Isolation and identification of bacterial strains with insecticidal activities from *Rhynchophorus ferrugineus* Oliver (Coleoptera: Curculionidae). *J. Appl. Entomol.* 140(8):617-626.
- Ramzan U, Abid K, Zafar MA, Anwar AM, Nadeem M, Tanveer U and Fatima U, 2023. *Trichoderma*: multitasking biocontrol agent. *Int. J. Agric. Biosci.* 12(2):77-82.
- Singh G, Sharma JR and Hoondal GS, 2008. Chitinase production by *Serratia marcescens* GG5. *Turk. J. Biol.* 32(4):231-236.
- Sundaravadivelan C and Padmanabhan MN, 2014. Effect of mycosynthesized silver nanoparticles from filtrate of *Trichoderma harzianum* against larvae and pupa of dengue vector *Aedes aegypti* L. *Environ. Sci. Pollut. Res.* 21:4624-4633.
- Trejo-López JA, Rangel-Vargas E, Gómez-Aldapa CA, Villagómez-Ibarra JR, Falfán-Cortes RN, Acevedo-Sandoval OA and Castro-Rosas J, 2022. Isolation and Molecular Identification of *Serratia* Strains Producing Chitinases, Glucanases, Cellulases, and Prodigiosin and Determination of Their Antifungal Effect against *Colletotrichum siamense* and *Alternaria alternata* In Vitro and on Mango Fruit. *Int. J. Plant Biol.* 13(3):281-297.
- Zhang CX, Yang SY, Xu MX, Sun J, Liu H, Liu JR, Liu H, Kan F, Sun J and Lai R, 2009. *Serratia nematodiphila* sp. nov., associated symbiotically with the entomopathogenic nematode *Heterorhabditis chongmingensis* (Rhabditida: Rhabditidae). *Int. J. Syst. Evol. Microbiol.* 59(7):1603-1608.
- Zhong S and Chen Y, 2012. Research progress in application of entomopathogenic fungi. *Fujian. Agric. Sci. Technol.* 43:58-61.
- Zhu H, Qin W, Huang S, Yan W and Sun X, 2010. Isolation and identification of an entomopathogenic fungus strain of *Rhynchophorus ferrugineus* Oliver. *Acta Phytopythologica Sinica.* 37(4):336-340.

